



Brush Up on Oral Health

July 2018

- **Sugar and Children's Oral Health**
 - **How Often Children Consume Foods and Drinks with Sugar During the Day Matters**
 - **Finding Hidden Sugar**
 - **Tips to Help Parents Limit Their Child's Sugar Intake**
- **Cook's Corner: Watermelon Stars**

Did You Know?

- **On average, children in the United States consume 19 teaspoons of added sugar per day.**
- **Children under age 2 should not be served any foods or drinks with added sugar.**
- **Children over age 2 should not be served foods or drinks with more than 6 teaspoons of added sugar per day.**



Sugar and Children's Oral Health

Children who consume foods and drinks containing natural and/or added sugar frequently (for example, every hour) during the day are more likely to develop tooth decay than those who consume them less often. Parents and Head Start staff may not know that many of the foods and drinks they give children contain sugar.

This issue of *Brush Up on Oral Health* is all about sugar. It describes how consuming foods and drinks containing sugar often during the day increases the risk of developing tooth decay. It shares information about how to read food labels to identify hidden sugars in products and tips for Head Start staff


to share with parents about limiting their child's sugar intake. It also provides a recipe for a healthy snack that can be made in a Head Start classroom or at home.

How Often Children Consume Foods and Drinks with Sugar During the Day Matters

Sugar plays a key role in tooth decay. (See the [April 2013](#) issue of *Brush Up on Oral Health* for more information on tooth decay.) Most foods, like milk and milk products, fruit, vegetables, grains, and processed and prepared foods, contain sugar.

Bacteria that cause tooth decay break down foods and drinks that contain sugar to form acid. Each time a person consumes foods or drinks containing sugar, acid is in the mouth for 20 to 40 minutes. Children who are fed meals and snacks at scheduled times are at lower risk for developing tooth decay than children who are fed often during the day.

The peaks in the top graph (Regularly Scheduled Meals and Snacks = Few Acid Attacks) show that acid is in the child's mouth five times during the day, for a total of a little more than 1½ hours. The peaks in the bottom graph (Fed Often During the Day = Many Acid Attacks) show that acid is in the child's mouth nine times during the day, for a total of 3 hours (about twice as long as for children who are fed meals and snacks at scheduled times). If a child consumes foods and drinks containing sugar often, over time the child is more likely to develop tooth decay.



Finding Hidden Sugar

Many foods and drinks contain added sugar. Sugar in foods can be listed by many different names. The best place to check for sugar is in the ingredients list on the food label. Look for words like:

- Beet sugar
- Brown sugar
- Cane sugar
- Corn sweeteners
- Corn syrup
- Cane juice
- High fructose corn syrup
- Honey
- Malt syrup
- Maple syrup
- Molasses
- Raw sugar
- White sugar

Ingredients: Whole Grain Oats (includes the oat bran), Sugar, Modified Corn Starch, Honey, Brown Sugar Syrup, Salt, Tripotassium Phosphate, Canola and/or Rice Bran Oil, Natural Almond Flavor. Vitamin E (mixed tocopherols) Added to Preserve Freshness.

Tips to Help Parents Limit Their Child's Sugar Intake

Head Start staff can help prevent tooth decay by teaching parents about hidden sugar in foods and drinks and about feeding and eating habits that can reduce their child's risk for developing tooth decay.

- Explain to parents the role sugar plays in the tooth decay process and that how often a child consumes foods and drinks containing sugar can make a big difference.
- Show parents how to identify sugars listed on ingredient labels.
- Work with parents to set up a schedule for serving meals and snacks. Encourage parents not to feed their child or graze on foods and drinks often through the day, especially those containing natural and/or added sugar.

Cook's Corner: Watermelon Stars

Here is a delicious healthy snack that children can make in a Head Start classroom or at home with their families.

Ingredients

- 1 small seedless watermelon
- 1 cup low-fat cottage cheese
- ½ cup blueberries

Directions

1. Cut the watermelon into ½-inch-thick slices, then use a star-shaped cookie cutter to cut the slices into stars.
2. Top each star with a teaspoonful of cottage cheese.
3. Place a blueberry on top of the cottage cheese.

Makes 10–12 servings

Safety tip: An adult should slice the ingredients.



Contact Us

The National Center on Early Childhood Health and Wellness welcomes your feedback on this issue, as well as your suggestions for topics for future issues. Please forward your comments to health@ecetta.info or call 866-763-6481.

Subscribe or view all issues of [Brush Up on Oral Health](#) on the Early Childhood Learning and Knowledge Center.

This issue was developed with funds from grant #U44MC30806 for the U.S. Department of Health and Human Services, Administration for Children and Families, Office of Head Start, Office of Child Care, and Health Resources and Services Administration, Maternal and Child Health Bureau, by the National Center on Early Childhood Health and Wellness. This resource may be duplicated for noncommercial uses without permission.

This publication is in the public domain, and no copyright can be claimed by persons or organizations.

School readiness begins with health!



ADMINISTRATION FOR
CHILDREN & FAMILIES



NATIONAL CENTER ON
Early Childhood Health and Wellness